

# RELIANCE CALF 2000



**RELIANCE CALF 2000 is a high protein, mineralised and vitamised calf feed designed to support calf health and vitality.**

## RELIANCE CALF 2000 CONTAINS:

- Provides high digestibility for young calves
- Contains GutBoosta™ with prebiotics and essential oils.
- Quality tested to meet stated nutrient specifications.
- Contains the coccidiostat Bovatec®.
- Made from selected quality palatable ingredients.
- Contains essential vitamins (including B vitamins) and minerals.

## FEEDING RECOMMENDATIONS

Feed RELIANCE CALF 2000 from the first week of life after good colostrum intake has been achieved. Offer fresh daily and clean out troughs regularly; hand feeding immediately after milk feeds can encourage early intake. Additional coccidiostat may be required in the milk until pellet intake is sufficient to aid in the control of coccidiosis.

Wean off milk gradually at a minimum 6 weeks of age and/or 65-80kg when consuming between 1 to 1.5kg RELIANCE CALF 2000 depending on the calf size and breed. Continue to feed RELIANCE CALF 2000 at 1-1.5kg/head/day for 1-2 months after weaning from milk, transitioning to 16% version if pasture quality permits.

Provide RELIANCE CALF 2000 along with whole milk or quality calf milk replacer and long forage such as grass or hay. Ensure calves have access to reachable drinking water at all times.

Do not feed to any animal other than those stipulated on the label without consulting a nutritionist.

Please ensure product is stored in a cool, dry and vermin free environment.

## INGREDIENTS SELECTED FROM

Barley, wheat, triticale, maize grain and grain by-products, peas, beans, soyabean meal, oilseed meals, molasses, vegetable oils, major minerals, trace minerals, vitamins A, D and E, range of B vitamins including Biotin, prebiotic, essential oils, coccidiostat (Bovatec® 20 CC), mould inhibitor, flavour. RELIANCE CALF 2000 does not contain palm kernel meal.

Additional copper or selenium should only be given following consultation with a nutritionist or veterinarian.

## TYPICAL ANALYSIS (APPROXIMATE ON A DRY MATTER BASIS)

Protein	20%
Moisture	13%
Fat	< 3%

**You can rely on Reliance**

**RELIANCE  
FEEDS**

## COCCIDIOSTAT

### WARNING STATEMENTS

READ LABEL BEFORE USE

FOR ANIMAL TREATMENT ONLY

KEEP OUT OF THE REACH OF CHILDREN

### Reliance Calf 2000 calf feed with 500g/tonne Bovatec® 20CC.

**CONTAINS** Bovatec® 20CC which is registered pursuant to the ACVM Act 1997, No. A009679.

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

**INDICATIONS** Contains 100mg/kg of lasalocid sodium as 500g/tonne Bovatec® 20CC as an aid in the control of coccidiosis caused by Eimeria species in cattle. Feed 0.5kg of Reliance Calf 2000 per 50kg calf to provide 1mg of lasalocid per kg of body weight. Ensure young animals have access to Reliance Calf 2000 at least 2-3 weeks prior to weaning as coccidial challenge is most likely to occur in the first eight weeks after weaning.

**DIRECTIONS FOR USE** This product must not be used for growth promotion in ruminants intended for human consumption. By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

**DOSE RATE** For coccidiosis control, minimum dose rate of 5mg Bovatec®/kg live weight (1mg lasalocid sodium/kg live weight) is required per day. Up to twice this dose rate is indicated where severe coccidial challenge is expected or a wide range of individual live weights occur within a group.

**CAUTION** Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec®, as ingestion by these species may be fatal.

Do not feed to deer, milking goats, sheep, laying poultry and other non-ruminant animals.

## CONTRAINDICATIONS

- Care must be exercised when feeding concurrently with other antimicrobials.
- Do not feed with other ionophores, e.g. monensin capsules, liquid or premix.
- Do not exceed the recommended dose rates.
- Do not feed more than a total of 400mg lasalocid sodium/head/day to lactating cattle.
- Not to be used for single-dose treatment.
- Disposal: avoid contamination of any water supply with this feed or its empty bag.

**WITHHOLDING PERIOD** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**Cattle**

**Meat:** Nil

**Milk:** Cattle Nil

**Not to be used for bobby calves**

## MANUFACTURED FOR AND DISTRIBUTED BY:

**Farmlands Nutrition**

Private Bag 1968, Dunedin 9054, New Zealand.